

# Tuning and Validating HYCOM's KPP Mixed-Layer

Alan J. Wallcraft and E. Joseph Metzger

Naval Research Laboratory

A. Birol Kara

COAPS, Florida State University

2003 Layered Ocean Model Users' Workshop

February 10, 2003

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>10 FEB 2003</b>		2. REPORT TYPE		3. DATES COVERED <b>00-00-2003 to 00-00-2003</b>	
4. TITLE AND SUBTITLE <b>Tuning and Validating HYCOM's KPP Mixed-Layer</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>Naval Research Laboratory,Stennis Space Center,MS,39529</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release; distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>Layered Ocean Modeling Workshop (LOM 2003), Miami, FL, Feb 2003</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>Same as Report (SAR)</b>	18. NUMBER OF PAGES <b>22</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			


- Long-term Global HYCOM Objective
  - To depict and predict the 3D global ocean state at fine resolution ( $0.08^\circ$  on the equator,  $\sim 7$  km at mid-latitudes)
  
- Strategy
  - Begin with  $0.72^\circ$  resolution global HYCOM and optimize the KPP mixed layer performance (the subject of this presentation)
  - Before the end of FY03, start running and development of  $0.24^\circ$  global HYCOM
  - Submitted DoD HPC Challenge proposal (FY04-06) to continue development at  $0.24^\circ$  and start  $0.08^\circ$  model development

## **0.72 degree Global Domain**


- Pan-Am Global Grid
  - 0.72 degree equatorial Mercator 78S-47N
  - Arctic bi-polar patch above 47N
    - \* Low resolution global had patch at 59N
    - \* Can't include Hudson Bay
  - Double latitudinal resolution near the equator
  - Halve latitudinal resolution in Antarctic
- Coastline at 50m isobath
  - Closed Bering Strait
  - No Sigma (terrain-following) vertical coordinate
- Same 26-layers as 0.08 degree Atlantic
- First Z-level 3 m, increases 1.125x up to 12 m



## 47N: SCPY (km)



## 47N: Grid Aspect Ratio (SCPX/SCPY)



## **0.72 degree Global Standard Configuration**

- KPP mixed layer
- Energy-Loan ice model
- Sigma-theta (some Sigma-2 runs, not discussed here)
- Horizontal diffusion chosen to suppress eddies
- Initialize from Levitus
- ECMWF Reanalysis monthly mean forcing
  - Plus 6-hrly wind anomalies from sep94-sep95
- Annual mean of 10 largest rivers via precip bogus
- Strong relaxation to monthly Levitus SSS
  - Necessary to prevent SSS drift
- Weak relaxation to monthly Levitus SST
  - Parameterizes SST's effect on longwave radiation


## Longwave Radiation and SST

- Longwave Radiation is sum of:
  - Upward blackbody longwave radiation
    - \*  $Q_{bb} = -0.98 (5.67 \times 10^{-8}) (T_s + 273.16)^4$
  - Downward atmospheric longwave flux
    - \* Highly dependent on cloudiness
    - \* Unknown dependence on SST (assume independent)
- If longwave was calculated using a SST of  $T_{so}$ :
  - $Q_{lw}(T_s) = Q_{lw}(T_{so}) + Q_{bb}(T_s) - Q_{bb}(T_{so})$
  - $Q_{lw}(T_s) = Q_{lw}(T_{so}) + Q'_{bb}(T_s - T_{so})$
  - $Q'_{bb} = -0.98 (5.67 \times 10^{-8}) 4 (T_s + 273.16)^3$
- Or, approximately (in  $W/m^2$ ):
  - $Q'_{bb} = -4.506 - 0.0554 T_s$
- So,  $-5.3 W/m^2/degC$  is a good longwave correction
  - Equivalent to 3.5 m reference thickness and 30 day e-folding SST relaxation
  - 10x weaker than typically SSS relaxation
- Ocean Model Intercomparison Project includes  $(Q_{bb}(T_s) - Q_{bb}(T_{so}))$  as a longwave correction

## Variable Turbidity

- Penetrating solar (shortwave) radiation is important for accurate SST
- HYCOM parameterizes turbidity via a “two band” version of Jerlov water types
  - One of 5 classes, red and blue bands
  - Same everywhere in time and space
- NLOM used “single band” kPAR turbidity
  - Photosynthetically Available Radiation
  - kPAR from SeaWIFS k490
  - monthly climatological fields
- Single band approach only OK for bulk mixed layer
- Added two band Jerlov-like kPAR scheme
  - kPAR isn't a good fit to Jerlov
  - Much better than single class everywhere
- Need a better scheme:
  - 3-band based on chlorophyll (Morel & Antoine, 1994)?
  - Jerlov-like based on k490?

## Annual SeaWiFS KPAR



## SST Metrics I

- Run for 5 years and form monthly means
  - May not be long enough
  - Length of run vs confidence in results
  - Takes two days on 64 IBM POWER4 cpus
- Compare monthly SST to Reynolds and Smith (R&S) climatology
  - Monthly anomalies
  - Annual mean difference
  - RMS difference
  - Correlation Coefficient
  - Skill Score
    - \* Correlation squared - Unconditional Bias - Conditional Bias
    - \* Maximum is 1, but minium is -infinity
    - \* Measure of error w.r.t. seasonal cycle (i.e. w.r.t. standard deviation)
    - \* Use a minimum of 1 degC for standard deviation
      - Still get poor skill scores near equator


## SST Metrics II


- Purpose of comparison is to find “good enough” configuration
  - Assume that “skill” on climatological forcing is maintained on interannual forcing
  - Is monthly thermal climatological forcing enough?
  - NLOM experience suggests that this is OK, but can’t be certain until we run interannually with HYCOM
- Targets:
  - Annual mean error  $< 0.5$  degC
  - Correlation Coefficient  $> 0.6$
  - Skill Score  $> 0.3$
- Use zonal averages to reduce amount of data
  - Average not necessarily best statistic
    - \* A few large negative skill scores can dominate the average
  - Same targets as for full field





## Simulation History

- Expt 1.5:
  - Standard KPP
  - Jerlov Ia everywhere
  - No rivers
  - No SST relaxation
- Expt 1.8:
  - Monthly kPAR turbidity
  - Annual rivers
- Expt 3.3:
  - “Longwave” SST relaxation
- Expt 3.9:
  - Modified KPP shear instability
- Expt 4.2:
  - Twin of 3.3 (same winds)
  - FNMOC thermal forcing (average 98-01)












## 03.3 vs R&S SST: Mean Error



## 04.2 vs R&S SST: Mean Error




## 03.3 vs R&S SST: Correlation






## 03.3 vs R&S SST: Skill Score



## 04.2 vs R&S SST: Skill Score



## Conclusions

- KPP is performing well in HYCOM
- Western Equatorial Atlantic a major trouble spot
  - Modifying KPP shear instability helps
  - More needs to be done
- In general, most of the SST error is in the annual mean
- Not yet clear how much is due to forcing and how much due to KPP